

## A double-hurdle approach to modelling tobacco consumption in Italy

Aristei, David; Pieroni, Luca

Postprint / Postprint

Zeitschriftenartikel / journal article

Zur Verfügung gestellt in Kooperation mit / provided in cooperation with:

[www.peerproject.eu](http://www.peerproject.eu)

### Empfohlene Zitierung / Suggested Citation:

Aristei, D., & Pieroni, L. (2008). A double-hurdle approach to modelling tobacco consumption in Italy. *Applied Economics*, 40(19), 2463-2476. <https://doi.org/10.1080/00036840600970229>

### Nutzungsbedingungen:

Dieser Text wird unter dem "PEER Licence Agreement zur Verfügung" gestellt. Nähere Auskünfte zum PEER-Projekt finden Sie hier: <http://www.peerproject.eu> Gewährt wird ein nicht exklusives, nicht übertragbares, persönliches und beschränktes Recht auf Nutzung dieses Dokuments. Dieses Dokument ist ausschließlich für den persönlichen, nicht-kommerziellen Gebrauch bestimmt. Auf sämtlichen Kopien dieses Dokuments müssen alle Urheberrechtshinweise und sonstigen Hinweise auf gesetzlichen Schutz beibehalten werden. Sie dürfen dieses Dokument nicht in irgendeiner Weise abändern, noch dürfen Sie dieses Dokument für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, aufführen, vertreiben oder anderweitig nutzen.

Mit der Verwendung dieses Dokuments erkennen Sie die Nutzungsbedingungen an.

**gesis**  
Leibniz-Institut  
für Sozialwissenschaften

### Terms of use:

This document is made available under the "PEER Licence Agreement". For more information regarding the PEER-project see: <http://www.peerproject.eu> This document is solely intended for your personal, non-commercial use. All of the copies of this documents must retain all copyright information and other information regarding legal protection. You are not allowed to alter this document in any way, to copy it for public or commercial purposes, to exhibit the document in public, to perform, distribute or otherwise use the document in public.

By using this particular document, you accept the above-stated conditions of use.

Mitglied der  
  
Leibniz-Gemeinschaft



### A Double-Hurdle Approach to Modelling Tobacco Consumption in Italy

Journal:	<i>Applied Economics</i>
Manuscript ID:	APE-06-0020.R1
Journal Selection:	Applied Economics
JEL Code:	C24 - Truncated and Censored Models < C2 - Econometric Methods: Single Equation Models < C - Mathematical and Quantitative Methods, D12 - Consumer Economics: Empirical Analysis < D1 - Household Behavior and Family Economics < D - Microeconomics
Keywords:	double-hurdle models, tobacco consumption, limited dependent variables, Box-Cox transformation

powered by ScholarOne  
Manuscript Central™

# A Double-Hurdle Approach to Modelling Tobacco Consumption in Italy

David ARISTEI<sup>1</sup> – Luca PIERONI<sup>2</sup>

## Abstract

*This paper analyses the determinants of tobacco expenditures for a sample of Italian households. A Box-Cox double-hurdle model adjusted for heteroscedasticity is estimated to account separate individual decisions concerning smoking participation and tobacco consumption and to correct for non-normality in the bivariate distribution of the error terms. Nested univariate and bivariate models are found to be excessively restrictive, supporting the adequacy of a generalized specification.*

*Estimation results show that consumption decisions are significantly affected by income and demographic characteristics. In particular, income positively impacts tobacco expenditure, while participation probability substantially declines as age increases. The existence of significant gender differences in both smoking participation and tobacco consumption patterns is found, while high education and white collar occupation reduce the likelihood to smoke and tobacco expenditure levels. Single adult households have a lower probability of smoking initiation even if, conditional on smoking, they consume more. Finally, complementarity between tobacco and alcohol beverages suggests the necessity of joint public health strategies.*

J.E.L. classification: C24, D12

*Keywords: tobacco consumption, double-hurdle models, limited dependent variables, Box-Cox transformation.*

---

<sup>1</sup> *Department of Economics, Finance and Statistics, University of Perugia and Department of Economic Sciences, University of Verona and.*

<sup>2</sup> *Department of Economics, Finance and Statistics, University of Perugia.*

**Corresponding author:** Luca Pieroni, Department of Economics, Finance and Statistics,  
University of Perugia, via Pascoli 20, 06123 Perugia;  
Tel. +390755855280, Fax +390755855299, e-mail: lpieroni@unipg.it

## 1. Introduction

In the last years the empirical literature has produced a large body of evidence on the price and non-price determinants of tobacco demand. One strand of literature has adopted an aggregate time series approach to provide empirical support to the rational addiction model proposed by Becker and Murphy (1988) (Chaloupka, 1991; Becker et al., 1994; Bask and Melkersson, 2004). On the other hand, the growing availability of microdata from household expenditure surveys has allowed to model tobacco consumption accounting for zero observations and simultaneously exploiting the richness of survey data information to control for heterogeneous individual (or household) behaviour (Jones, 1989, 1992; Blaylock and Blisard, 1992; Garcia and Labeaga, 1996; Yen, 2005a). From a policy perspective, cross-sectional surveys enables to improve the knowledge of the impacts of socio-demographic variables on tobacco expenditure and help the design of public health programs to achieve smoking-reduction objectives.

While it would be interesting to obtain simultaneous empirical responses concerning addiction, censoring and heterogeneity in tobacco consumption decisions in Italy, the absence of a true panel data does not enable us to account for addictive behaviours while controlling for demographic and socio-economic characteristics<sup>1</sup>. Thus, in this paper we investigate household tobacco expenditures, addressing the issues connected to limited dependent variable models by an approach based on a double-hurdle specification (Cragg, 1971; Jones, 1989; Yen and Jones, 1996; Su and Yen, 2000). Several empirical studies (Blundell and Meghir, 1987; Blaylock and Blisard, 1993;

---

<sup>1</sup> Only few countries give a panel data structure to their household expenditure surveys. Recently, Labeaga (1999) and Jones and Labeaga (2003), using a panel of Spanish households (the Continuous Family Expenditure Survey) have attempted to test rational addiction and simultaneously account for censoring and unobservable heterogeneity.

1  
2  
3  
4 Garcia and Labeaga, 1996; Yen and Jones, 1996) have shown the inadequacy of the  
5  
6 standard Tobit model (Tobin, 1958) in cross-sectional analysis of tobacco consumption,  
7  
8 connected with its failure in accounting for differences concerning the generation of  
9  
10 zero observations. Since it is implausible that all zero observations in tobacco  
11  
12 consumption arise from standard corner solutions generated by a constrained budget, we  
13  
14 emphasize the importance of a double-hurdle specification distinguishing between  
15  
16 abstentions and corner solutions.  
17  
18  
19

20  
21 As shown by Arabmazard and Schmidt (1982), maximum likelihood estimates are  
22  
23 sensitive to misspecification issues and they will be inconsistent if the bivariate  
24  
25 normality assumption is violated. One way to correct for the non-normality of the error  
26  
27 terms, providing to generalize the standard double-hurdle model, consists in applying a  
28  
29 Box-Cox transformation on the dependent variable (Yen, 1993; Yen and Jones, 2000).  
30  
31 In this way we can encompass, besides the standard double-hurdle model, a wide range  
32  
33 of specifications, that differ for the distributional assumptions on the error terms, and  
34  
35 test the best model to rationalize the data.  
36  
37  
38  
39

40  
41 Standard Tobit model, standard Double-hurdle model with independence and Box-  
42  
43 Cox Tobit model seem to be restrictive specifications, confirming both the violation of  
44  
45 the bivariate normality assumption and the existence of separate individual decisions on  
46  
47 participation and consumption. Moreover, the independence of the errors for the Box-  
48  
49 Cox double-hurdle is not rejected by the data, concluding that the non-normal  
50  
51 specification with independent errors is the best model to account for household  
52  
53 behaviours on tobacco expenditures.  
54  
55

56  
57 The remainder of the paper is organized as follows: the next section outlines the  
58  
59 theoretical framework upon which the empirical models are based. In Section 3 we  
60

1  
2  
3  
4 discuss econometric methods. In particular, both the standard double-hurdle model and  
5  
6 the extended specification based on the Box-Cox transformation are presented. In  
7  
8 Section 4 data used in the empirical analysis, taken from 2002 Italian Household Budget  
9  
10 Survey, are discussed and factors influencing participation and consumption equations  
11  
12 are examined. In Section 5 specification and estimation results are presented and  
13  
14 discussed, with specific attention devoted to the nested strategy used to derive the best  
15  
16 specification and to the analysis of the estimated parameters and elasticities. Section 6  
17  
18 offers some concluding comments.  
19  
20  
21  
22  
23  
24  
25

## 26 **2. Theoretical Framework**

27  
28 Standard consumer choice theory assumes that every individual (or household) is a  
29  
30 potential consumer of all goods. However, for commodities like tobacco this may not be  
31  
32 true and, regardless of price and income levels, some individuals cannot be induced to  
33  
34 smoke. In these cases, zero observations are not the result of economic non-  
35  
36 consumption, but they may be determined by other behavioural factors apart from prices  
37  
38 and income. Following Pudney (1989), alternative models of tobacco demand have been  
39  
40 derived by using discrete random preferences regimes. According to this approach,  
41  
42 smokers are hypothesized to have a different preference structure than non-smokers.  
43  
44 Observed zero expenditure then reflect either the decision to not smoke or a standard  
45  
46 corner solution and hence only potential smokers determine the parameters of tobacco  
47  
48 Engle curve (Blaylock and Blisard, 1993).  
49  
50  
51  
52  
53

54 Given this setting, individual's utility function takes the following form:

$$55 U = U(dc_1, c_2, \dots, c_n; w) \quad (1)$$

56  
57  
58  
59  
60

where  $c_1$  is the quantity of tobacco (with price  $p_1$ ),  $c_2, \dots, c_n$  represent all other goods,  $w$  is a vector of demographic variables representing the qualitative characteristics of smoking, and  $d$  is binary variable which is equal to one if an individual is an actual or potential smoker and zero otherwise. If  $d$  always equals one, every individual is assumed to be a potential smoker and observed zero consumption is a standard corner solution.

Equation (1) can be alternatively rewritten as:

$$U = dU^s(c_1, c_2, \dots, c_n; w) + (1-d)U^{ns}(c_2, \dots, c_n; w) \quad (2)$$

where  $U^s$  is the utility function of smokers (actual and potential) and  $U^{ns}$  for non-smokers. For a non-smoker, given that  $c_1$  does not enter  $U^{ns}(c_2, \dots, c_n; w)$  and  $p_1$  is in any case positive, the optimal tobacco consumption level is  $c_1^* = 0$ . For actual and potential smokers, the optimal level of  $c_1$  is determined by solving the following constrained utility maximization:

$$\max_{c_1, \dots, c_n} \{U^s(c_1, c_2, \dots, c_n; w)\} \quad \text{s.t.} \quad p'c = m \quad (3)$$

where  $p$  is a vector of prices (including  $p_1$ ) and  $m$  is individual's (or household's) budget. Assuming the utility function  $U^s(c_1, c_2, \dots, c_n; w)$  to be continuous, increasing, and quasi-concave, then the notional demand for tobacco can be expressed as a demand function  $f(p, m; w)$  and the corresponding expenditure equation can be denoted as  $g(m; w)$ . As price information was not collected in the ISTAT Italian Household Budget Survey, we assume that all households face the same relative prices (Yen and Jensen, 1996). The notional demand and expenditure have been derived as the results of utility maximization with only the budget constraint, given individual (or household)

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

characteristics. However, as the quantity and expenditure are also subject to a non-negativity constraint, the optimal level of tobacco expenditure ( $e_{c_1}^*$ ) can be either an interior solution or a corner solution (that is:  $e_{c_1}^* = \max\{0, g(m;w)\}$ ), justifying the use of a double-hurdle specification for modelling tobacco consumption, since it jointly accounts for both abstentions and corner solutions.

In the next section, alternative empirical models are developed in order to account for the presence of different zero-generating mechanisms in tobacco consumption.

### 3. Econometric Specification

Recent empirical analyses have shown the inadequacy of the standard Tobit model in cross-sectional analysis of tobacco consumption, stressing the relevance of a double-hurdle approach for microeconomic analysis of tobacco consumption (Jones, 1989; Blaylock and Blisard, 1992; Garcia and Labeaga, 1996; Yen and Jones, 1996; Labeaga, 1999). The main feature of the double-hurdle model is that participation and consumption decisions are assumed to stem from two separate individual choices and the determinants of the two decisions are allowed to differ.

The double-hurdle model, originally proposed by Cragg (1971), assumes that two separate hurdles must be passed before a positive level of consumption can be observed. In the context of tobacco consumption analysis, the first hurdle involves the decision of whether or not to smoke (*participation decision*). It is reasonable to assume that the choice of smoking is not only an economic decision, but also influenced by social and demographic factors which are independent of the quantity consumed. The second hurdle concerns the level of tobacco consumption to choose (*consumption decision*).



We now turn to the econometric specification of the model. Both hurdles are assumed to be linear in the parameters  $(\alpha, \beta)$ , with disturbance terms  $u$  and  $v$  randomly distributed with a bivariate normal distribution. The matrices  $z$  and  $x$  include the variables that are assumed to influence participation and consumption decisions, respectively. Formally, following Jones (1989) and Pudney (1989), the bivariate model can be written as:

i) *Observed consumption:*

$$y_i = d \cdot y_i^{**} \quad (4)$$

ii) *Participation equation:*

$$w_i = z_i' \alpha + u_i \quad u_i \sim N(0, 1) \quad (5)$$

$$d = \begin{cases} 1 & \text{if } w > 0 \\ 0 & \text{otherwise} \end{cases}$$

iii) *Consumption equation:*

$$y_i^* = x_i' \beta + v_i, \quad v_i \sim N(0, \sigma^2) \quad (6)$$

$$y_i^{**} = \begin{cases} y_i^* & \text{if } y_i^* > 0 \\ 0 & \text{otherwise} \end{cases}$$

A positive level of tobacco consumption  $y$  is observed only if the individual (or the household) is a potential smoker ( $d = 1$ ) and actually consumes tobacco ( $y^{**}$ ). For this reason, in double-hurdle models, differently from Heckman selection model (Heckman, 1979), in which zeros are not affected by the consumption decision, observed zero expenditures are the result of either participation or consumption decisions and potential smokers may have zero tobacco expenditure.

In the present analysis, different double-hurdle models are used to analyze household tobacco expenditure patterns, with particular attention devoted to the specification of the stochastic structure of the model by testing whether the assumptions of bivariate

normality, homoscedasticity and independence of the error terms across participation and consumption equations are acceptable.

To account for the presence of heteroscedasticity, the variance of the error terms specified it as a function of a set of continuous variables:

$$\sigma_i = \exp(z_i' h) \quad (7)$$

where  $z_i$  is a vector of continuous variables included in  $x_i$  ( $z_i \in x_i$ ) and  $h$  is a conformable vector of coefficients (Yen, 1993; Yen and Jensen, 1996; Newman et al., 2003).

The assumption of correlated error terms allows for the possibility that participation and consumption decisions are simultaneously taken. In particular, we assume that  $u$  and  $v$  are distributed as a bivariate normal:

$$(u, v) \square BVN(0, \Sigma), \quad \Sigma = \begin{bmatrix} 1 & \sigma\rho \\ \sigma\rho & \sigma^2 \end{bmatrix} \quad (8)$$

where  $\rho$  is the correlation coefficient.

Denoting zero consumption as 0 and positive consumption as +, the likelihood function for the full double-hurdle model with heteroscedasticity correction and dependence between  $u$  and  $v$  can be written as:

*a) Heteroscedastic Double-hurdle model with dependent error terms*

$$L = \prod_0 [1 - \Phi(z_i' \alpha, x_i' \beta, \rho)] \prod_+ \left[ \Phi \left( \left[ z_i' \alpha + \frac{\rho}{\sigma_i} (y_i - x_i' \beta) \right] / \sqrt{1 - \rho^2} \right) \frac{1}{\sigma_i} \phi((y_i - x_i' \beta) / \sigma_i) \right] \quad (9)$$

where  $\Phi$  denotes the standard normal CDF (univariate or multivariate) and  $\phi$  is the univariate standard normal PDF.

In empirical applications it is common to assume that  $u$  and  $v$  are independent (i.e.  $\rho = 0$ ).

Under this hypothesis we obtain the original formulation proposed by Cragg (1971).

b) *Heteroscedastic Double-hurdle model with independent error terms*

$$L = \prod_0 [1 - \Phi(z'_i \alpha) \Phi(x'_i \beta) / \sigma_i] \prod_+ \left[ \Phi(z'_i \alpha) \frac{1}{\sigma_i} \phi((y_i - x'_i \beta) / \sigma_i) \right] \quad (10)$$

This model hypothesizes that the participation and consumption decisions are made separately and that there is a feedback effect from the level of consumption to the participation decision (Atkinson et al., 1984; Deaton and Irish, 1984; Blaylock and Blisard, 1993). In the present analysis we do not make any *a priori* assumption on the correlation structure of the error terms; differently from other studies (Yen and Jensen, 1996; Moffatt, 2005; Newman et al., 2003), the independence of error terms is not assumed as a maintained hypothesis, but its significance is tested by means of a Wald test. Moreover, it is worth noticing that the standard Tobit model is a nested version of the Cragg model, when  $\alpha_j = 0$  for  $j \neq 0$  and  $\alpha_0 = \infty$ , that is when  $p(u > -z'_i \alpha) = 0$  (i.e.  $\Phi(z'_i \alpha) = 1$ ), and a likelihood ratio test can be used to distinguish between the Tobit and the independent double-hurdle model.

A limitation of the standard double-hurdle specification is that it is built on the assumption of bivariate normality of the error terms. If the normality assumption is violated the maximum likelihood estimates of the model will be inconsistent. This may be particularly relevant when the model is applied to a dependent variable with a highly skewed distribution, as is often the case with survey data on tobacco expenditures.

As suggested by Yen (1993) and Jones and Yen (2000), one way to correct for the non-normality of the error terms consists in applying a Box-Cox transformation to the dependent variable, which gives:

$$y_i^T = \frac{y_i^\lambda - 1}{\lambda}, \quad \text{with } 0 < \lambda \leq 1 \quad (11)$$

where  $\lambda$  is an unknown parameter. It is worth noticing that the Box-Cox transformation includes, as special cases, a straightforward linear transformation ( $\lambda = 1$ ) and the logarithmic transformation ( $\lambda \rightarrow 0$ ), but normally we would expect the parameter  $\lambda$  to lie between these two extremes.

The Box-Cox double-hurdle model implies the following relationship between the transformed dependent variable and the latent variables,  $w$  and  $y^*$ :

$$y_i^T = \begin{cases} y_i^* & \text{if } y_i^* > -1/\lambda \text{ and } w_i > 0 \\ 0 & \text{otherwise} \end{cases} \quad (12)$$

where  $w_i$  and  $y_i^*$  are defined as in (7) and (8).

This specification relaxes the normality assumption on the conditional distribution of  $y_i$  and still allows stochastic dependence between the error terms of participation and consumption equations. The likelihood functions for dependent and independent double-hurdle models with heteroscedasticity correction can be written as:

c) *Box-Cox Heteroscedastic Double-hurdle model with dependent error terms*

$$L = \prod_0 \left[ 1 - \Phi \left( z_i' \alpha, \frac{x_i' \beta + 1/\lambda}{\sigma_i}, \rho \right) \right] \times \prod_+ \left[ \Phi \left( \left\{ z_i' \alpha + \frac{\rho}{\sigma_i} [(y_i^\lambda - 1)/\lambda - x_i' \beta] \right\} / \sqrt{1 - \rho^2} \right) y_i^{(\lambda-1)} \frac{1}{\sigma_i} \phi \left( [(y_i^\lambda - 1)/\lambda - x_i' \beta] / \sigma_i \right) \right] \quad (13)$$

d) *Box-Cox Heteroscedastic Double-hurdle model with independent error terms*

$$L = \prod_0 \left[ 1 - \Phi(z_i' \alpha) \Phi((x_i' \beta + 1/\lambda)/\sigma_i) \right] \prod_+ \left[ \Phi(z_i' \alpha) y_i^{(\lambda-1)} \frac{1}{\sigma_i} \phi \left( [(y_i^\lambda - 1)/\lambda - x_i' \beta] / \sigma_i \right) \right] \quad (14)$$

Model (c) is a general model that nests all the other three double-hurdle specifications presented and also encompasses a wide range of standard limited dependent variable models (Jones and Yen, 2000). Figure 1 summarizes the relationships between all the

models considered, showing the relevant restrictions on the likelihood function (13) implied by the nested specifications.

(Figure 1 about here)

The economic interpretation of limited dependent variable models frequently focuses on the analysis of the marginal effects of regressors on the expected value of  $y_i$  (Jones and Yen, 2000), which can be decomposed into an effect on the probability of purchase and an effect on the conditional level of expenditure<sup>2</sup>. The unconditional mean of  $y_i$  in the Box-Cox Double-hurdle model can be written as:

$$E(y_i) = P(y_i > 0)E(y_i | y_i > 0) \quad (15)$$

The conditional expectation of  $y_i$  is:

$$E(y_i | y_i > 0) = E(y_i | w_i > -z_i'\alpha, y_i^* > -x_i'\beta - \frac{1}{\lambda}) \quad (16)$$

and, assuming independence between error terms of participation and consumption<sup>3</sup>, can be written as:

$$E(y_i | y_i > 0) = \left[ \Phi \left( \frac{x_i'\beta + 1/\lambda}{\sigma_i} \right) \right]^{-1} \int_0^{\infty} \frac{y_i^\lambda}{\sigma_i} \phi \left( \frac{y_i - x_i'\beta}{\sigma_i} \right) dy_i \quad (17)$$

Given independence, the probability of a positive consumption level is:

$$P(y_i > 0) = \Phi(w_i'\alpha) \Phi \left( \frac{x_i'\beta + 1/\lambda}{\sigma_i} \right) \quad (18)$$

<sup>2</sup> This decomposition follows the approach proposed by McDonald and Moffitt (1980) for the decomposition of the unconditional mean of the dependent variable in the Tobit model.

<sup>3</sup> Here, for simplicity, we focus on the Independent Box-Cox Double-Hurdle model. Details on the derivation of the conditional mean for the Box-Cox Double-Hurdle model with dependent errors can be found in Jones and Yen (2000).

Marginal effects can be obtained by differentiating equations (16), (17) and (18) with respect to each explanatory variable<sup>4</sup>. From these marginal effects, elasticities can be derived. In particular, using equation (15), the elasticity of the conditional mean with respect to regressor  $x_{ij}$  can be written as:

$$e_j = \frac{\partial E(y_i)}{\partial x_{ij}} \frac{x_{ij}}{E(y_i)} = \frac{\partial P(y_i > 0)}{\partial x_{ij}} \frac{x_{ij}}{P(y_i > 0)} + \frac{\partial E(y_i | y_i > 0)}{\partial x_{ij}} \frac{x_{ij}}{E(y_i | y_i > 0)} \quad (19)$$

where the two addends are the elasticity of the probability of observing a positive expenditure ( $e_j^p$ ) and the elasticity of conditional consumption ( $e_j^{cc}$ ). For continuous variables, the elasticities are computed at the sample means. For categorical explanatory variables, marginal effects are used to compute percentage changes in probability, conditional level and unconditional level when the value of the variable shifts from zero to one, holding all the other variables constant (Yen and Jones, 1996; Newman et al., 2003).

#### 4. Data and Variables

The data used in the empirical analysis are taken from the 2002 Italian Household Budget Survey (IHBS), which is conducted by the Italian Central Statistics Office (ISTAT). This survey, together with Bank of Italy's Survey of Household Income and Wealth, represents the main and most comprehensive source of microdata for analysing consumption behaviours of Italian households. The ISTAT survey covers a random sample of 27499 households throughout the country and provides detailed information on family expenditures (non-durable and durable) as well as on household socio-economic and demographic characteristics. Data on non-durable consumption are collected in a diary that records household expenditures on a wide range of non-durable

<sup>4</sup> Analytical details on the derivation of conditional and unconditional marginal effects for the Box-Cox double-hurdle model can be found in Yen (1993) and Jones and Yen (2000).

1  
2  
3  
4 goods and services over a one-week period and is subsequently expressed on a monthly  
5  
6 basis. No price or quantity information is reported and consumption of each good or  
7  
8 service is measured as reported expenditure. Moreover, data refer to the household  
9  
10 rather than to individuals, providing no information on intra-household income  
11  
12 allocation and making it necessary to account for family composition effects when  
13  
14 analysing the demand for certain non-durable goods.  
15  
16  
17

18  
19 In this analysis, household monthly expenditure on cigarettes and tobacco is used as  
20  
21 the dependent variable. As already underlined, the ISTAT survey only records the value  
22  
23 of household expenditures and so quantity and quality effects cannot be identified and  
24  
25 analyzed. Since the survey does not provide any information on the presence of smokers  
26  
27 in the household, inference on participation in tobacco consumption has to be made on  
28  
29 the basis of recorded expenditure, identifying non-smoking households as those with  
30  
31 zero tobacco expenditure. Expenditures on tobacco, as all expenditures on commonly  
32  
33 used non durable goods, are monitored for only one week. The fact that the collection of  
34  
35 information only covers a seven-day period, with a 33.4 percent of consuming  
36  
37 households, introduces some room for undetected infrequency of purchases. However,  
38  
39 for commodities like tobacco, it is unlikely that observed zero expenditures capture  
40  
41 infrequent purchases and they probably stem from abstentions or corner solutions  
42  
43 (Garcia and Labeaga, 1996). Using household data to model tobacco consumption,  
44  
45 which is mainly the result of an individual decision, may generate some bias in the  
46  
47 analysis of consumption behaviours. For this reason, it is necessary to correctly account  
48  
49 for family size and composition effects as well as for the socio-economic characteristics  
50  
51 of the household, since the household situation plays an active role in modifying  
52  
53 individual preferences.  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4 Household tobacco expenditures are assumed to be expressible as a linear  
5 combination of explanatory variables that are assumed to separately affect participation  
6 and consumption decisions. The explanatory variables selected, together with their  
7 sample statistics, are presented and fully described in Table 1. They include household  
8 income (proxied by total expenditure), alcohol consumption, the head of the  
9 household's age, and dummy variables indicating education, marital status, gender and  
10 working status of the household's head, the percentage of adult male members, the  
11 presence of children under fourteen years old, home ownership, and participation in  
12 alcohol consumption.  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24

25  
26 All expenditure variables are expressed in real terms, by deflating current values  
27 using ISTAT regional price indexes, in order to account for price variability across  
28 regions. Moreover, in order to obtain a per-equivalent adult measure of household  
29 consumption, all expenditure variables are adjusted for family size by using the  
30 modified OECD equivalence scale, which assigns a weight equal to one to the first adult  
31 in the household, 0.5 to each other adult and 0.3 to each child under fourteen years of age.  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45

*(Table 1 about here)*

## 46 **5. Results**

47  
48  
49 In this section, estimation results are presented and discussed, with specific attention  
50 devoted firstly to the choice of the most appropriate model specification and then to the  
51 analysis of the effects of explanatory variables by calculating and decomposing  
52 elasticities.  
53  
54  
55  
56  
57  
58  
59  
60



### 5.1. Model Specification

One of the main objective of this paper is to test whether univariate or bivariate models are adequate for analysing tobacco consumption behaviour of Italian households.

All the double-hurdle specifications discussed in Section 3 have been estimated by maximizing the logarithm of the likelihood functions (9), (10), (13) and (14). One parameter estimation issue in double-hurdle models concerns the choice of the regressors for participation and consumptions equations. As it is known, the choice of the explanatory variables to be included in the two hurdle does not rest on any a priori theory and may be somewhat arbitrary. Given that the inclusion of the same set of regressors in each hurdle makes the parameters identification difficult, exclusion restrictions must be imposed<sup>5</sup>. In empirical applications the first hurdle is usually assumed to be a function of non-economic factors affecting household's smoking decision, so that economic variables can be excluded from the first equation (Newman et al, 2003). Their exclusion is motivated by the discrete random preference theory, according to which sample selection is determined exclusively by non-economic factors (Pudney, 1989; Yen, 2005a).

The foregoing arguments require, before presenting estimation results, a discussion of the explanatory variables included in the model. The independent variables considered are intended to encompass the determinants of both smoking participation and tobacco consumption decisions and their choice rests on suggestions taken from previous empirical literature (Jones, 1989; Blaylock and Blisard, 1992; Garcia and Labeaga, 1996;

---

<sup>5</sup> In estimating the final model we started with a specification that included all explanatory variables in both hurdles; insignificant variables were gradually dropped, with exclusion restrictions giving identification higher reliability.

1  
2  
3  
4 Yen and Jones, 1996; Yen, 2005a) and on more specific issues connected with  
5  
6 identification problems. Their inclusion in either participation or consumption equations  
7  
8 is justified by economic, demographic and sociological factor, such as habit formation,  
9  
10 information and social awareness on damages and health risks connected with smoking  
11  
12 and restriction to smoking due to security reasons or to health guidelines for public places.  
13  
14

15  
16 In this study, both participation and consumption decisions are postulated to be  
17  
18 influenced by the presence of children under fourteen years old within the household  
19  
20 (*Child014*), the percentage of adult male members (*PercMale*), age (*Age*), gender  
21  
22 (*MaleHH*), marital status (*Single*), education level (*Highedu*) and occupation  
23  
24 (*Whitecollar*) of the household's head.  
25  
26

27  
28 A dummy variable indicating the presence of children is included in the model  
29  
30 postulating that the household would not smoke or at least would attempt to moderate  
31  
32 tobacco consumption when small children are present (Blaylock and Blisard, 1993; Kerr  
33  
34 et al., 2004). The percentage of adult male members in the household (*PercMale*) is  
35  
36 included to account for gender differences in tobacco consumption and to test for the  
37  
38 presence of gender-differentiated smoking habits (Angulo et al., 2001). The education  
39  
40 variable (equal to one if the household's head has at least a high school education, zero  
41  
42 otherwise) has been often used in previous empirical studies (Jones, 1989; Blaylock and  
43  
44 Blisard, 1992, 1993; Yen, 2005b; Yen and Jensen, 1996; Garcia and Labeaga, 1996;  
45  
46 Yen and Jones, 1996), suggesting that individuals with better education may be more  
47  
48 aware of the health risks connected with consumption of unhealthy goods such as  
49  
50 tobacco and alcohol. Being a white-collar worker reflects the individual's social class  
51  
52 and may help in explaining how smoking habits vary among different social groups.  
53  
54 Age is considered to assess how age-related health problems affect smoking behaviours  
55  
56  
57  
58  
59  
60

1  
2  
3  
4 and to verify the existence of a significant lifecycle pattern for both tobacco  
5 participation and consumption decisions<sup>6</sup>.  
6  
7

8  
9 In the participation equation, we include an additional binary variable indicating  
10 whether the household displays a high expenditure level (over the 75<sup>th</sup> percentile of the  
11 observed distribution) on alcoholic beverages (*HighAlc*), as a proxy for habit formation  
12 tendencies (Blaylock and Blisard, 1993).  
13  
14  
15  
16  
17

18  
19 Specific variables accounting for economic conditions have been introduced in  
20 consumption equation. Total household expenditure (*Income*) is included as a proxy for  
21 current income. A variable indicating whether the household lives in a home that is  
22 owned or being bought (*OwnerOcc*) is included, following the suggestions of Atkinson  
23 et al (1984) and Jones (1989), as a proxy for wealth and economic stability. Further,  
24 household alcohol expenditure (*Alcohol*) is included as a proxy to verify the presence of  
25 complementary relationships with household expenditures on alcoholic beverages. The  
26 consumption equation also includes quadratic terms of age and income to capture  
27 possible non-linear relationships with tobacco expenditure (Jones, 1989; Garcia and  
28 Labeaga, 1996).  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44

## 45 *5.2 Statistical Tests and Estimation Results*

46  
47 In order to correctly analyze the determinants of tobacco expenditures and to model  
48 household smoking behaviour, one first task relates to the choice of the most  
49 appropriate specification. Our selection strategy consists in testing the bivariate model  
50 with dependent error terms, which is the most general specification and encompasses all  
51  
52  
53  
54  
55  
56  
57

---

58  
59 <sup>6</sup> Jones (1989) included the individual's age and its square as explicative variables, while Yen and Jensen  
60 (1996) used both household age composition and the age of the household head, showing significant life-  
cycle patterns for both participation and consumption decisions.

1  
2  
3  
4 the other bivariate and univariate models discussed in Section 3, against its nested  
5 alternatives, by means of conventional and adjusted (Vuong, 1989) likelihood ratio  
6 tests. However, it should be underlined that the validity of the LR tests strongly rests on  
7 the assumption that the general model is not misspecified (Yen and Jones, 1996); in  
8 particular homoscedasticity and normality assumptions should not be violated.  
9 Distributional assumptions assume crucial relevance in limited dependent variable  
10 models, since maximum-likelihood estimation will lead to inconsistent parameter  
11 estimates when normality and homoscedasticity are not fulfilled (Maddala and Nelson,  
12 1975; Arabmazard and Schmidt, 1982). For these reasons, preliminary tests for the  
13 validity of the distributional assumptions are necessary. To this end LR test for  
14 homoscedasticity and Pagan and Vella's (1989) moment base test for normality have  
15 been carried out on both Tobit and double-hurdle specifications<sup>7</sup>; the results are  
16 presented in Table 2.  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41

*(Table 2 about here)*

42 As can be noted, all equations present severe problems of non-normality and  
43 heteroscedasticity, with LR test values well above the relevant critical values in both  
44 Tobit and double-hurdle models. The violation of homoscedasticity requires allowance  
45 for heteroscedastic error terms in the univariate and bivariate specifications. Following  
46 Yen (1993), we relax homoscedasticity assumption by specifying standard deviation  $\sigma_i^2$   
47 as a function of the continuous variables of the model, as in equation (7), and allowing  
48 it to vary across observations. For this reason, all the models considered in the  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59

---

60 <sup>7</sup> Details on distributional tests in censored and limited dependent variable models can be found in Bera, Jarque and Lee (1984), Pagan and Vella (1989) and Wells (2003)

1  
2  
3  
4 remainder of the discussion account for heteroscedasticity, with a variance equation that  
5  
6 includes only the continuous regressors that result statistically significant in generating  
7  
8 heteroscedasticity<sup>8</sup>. Results of normality tests reveal that estimation of standard Tobit  
9  
10 and double-hurdle models may lead to inconsistent results, supporting the necessity of a  
11  
12 non-normal generalization of these models. As previously introduced in Section 3,  
13  
14 following Yen (1993) and Yen and Jones (1996, 2000), we consider a Box-Cox  
15  
16 transformation of the dependent variable that relaxes normality assumption on the  
17  
18 conditional distribution of  $y_i$  and includes as special cases linear and logarithmic  
19  
20 transformations. The results of the normality tests also can be interpreted as a strong  
21  
22 indication of the superiority of the univariate and bivariate Box-Cox generalizations  
23  
24 with respect to their standard counterparts.  
25  
26  
27  
28  
29

30  
31 Once the diagnostics of the model have been analyzed, we now turn to the choice of  
32  
33 the most appropriate model. As previously shown in Figure 1, all restricted models can  
34  
35 be obtained by placing the relevant restrictions on the likelihood function (13) and can  
36  
37 be interpreted as special cases of the Box-Cox double-hurdle model with dependent  
38  
39 error terms<sup>9</sup>.  
40  
41

42  
43 The specification tests carried out are reported in Table 3. Firstly, we tested the  
44  
45 hypothesis of independent errors between participation and consumption equations; the  
46  
47 issue of dependency in double-hurdle models is a problem of great relevance, but it has  
48  
49 often been disregarded in previous empirical works (Newman et al, 2003; Moffatt,  
50  
51  
52

---

53  
54 <sup>8</sup> In principle all explanatory variables can be included in the heteroscedasticity specification; however,  
55  
56 doing so would considerably increase the number of parameters to be estimated. So we focused our  
57  
58 attention only on the variables that are more likely to cause heteroscedasticity and then we tested  
59  
60 alternative specification excluding those variables that are not significantly different from zero.

<sup>9</sup> The Heckman sample selection model can also be obtained as a restricted specification, assuming that participation decision dominates consumption decision. Vuong specification test for non-nested models supports the inadequacy of the Heckman model. The results are not presented here, but they are available from the authors.

2005). The results of the LR test ( $\chi^2_{(1)} = 0.78$  with a p-value equal to 0.377) clearly indicates that dependency is not relevant; this result is in line with findings of Jones (1989), Blaylock and Blisard (1993) and Garcia and Labeaga (1996) and demonstrates that the independent Box-Cox double-hurdle model is an acceptable alternative to the dependent model<sup>10</sup>. On the basis of the results of Vuong specification test for nested models (Vuong, 1989), all the other restricted specifications are rejected, each with a *p-value* of less than 0.0001. The interpretation of these results is twofold. Firstly, they suggest the inadequacy of the univariate Tobit specification in modelling tobacco consumption behaviours, given the existence of separate participation and consumption decisions. On the other hand, the results give further support to the generalized specification to account for non-normal and heteroscedastic error terms. Thus, the model that best rationalizes tobacco expenditure data is the independent Box-Cox double-hurdle model.

*(Table 3 about here)*

Maximum-likelihood estimates are presented in Table 4. In order to account for differences in estimated parameters, we report the results of both standard and Box-Cox independent double-hurdle models, even if the discussion is focused only on the latter.

Analyzing the estimated parameters, it is possible to highlight that all the coefficients, with the exception of that of education in the consumption equation and that of occupational status in the participation equation, are significant at the one

---

<sup>10</sup> Smith (2003) puts into question the relevance of the dependent double-hurdle model itself, asserting that this model contains too little statistical information to support estimation of dependency, even when dependency is truly present.

1  
2  
3  
4 percent level. Before to show the effects of explanatory variables, we underline that the  
5  
6 Box-Cox parameter ( $\lambda = 0.1864$ ) is significantly different from one and zero; this  
7  
8 implies a Box-Cox transformation that is different from both a linear transformation and  
9  
10 a logarithm specification.  
11  
12

13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
*(Table 4 about here)*

Turning to explanatory variables, income has a positive effect on household tobacco consumption, while income squared, which accounts for non-linearity relationships, is negative. These outcomes are in line with the findings of Garcia and Labeaga (1996) and imply that tobacco expenditure rises as household income increases, but at a decreasing rate. The estimated age effects are negative and significant in both participation and consumption equations, indicating the existence of a strong lifecycle pattern (Kerr et al., 2004; Aristei et al., 2005). As in Yen and Jones (1996) and Yen (2005b), the probability of smoking and tobacco expenditure levels decrease with the age; moreover, the estimated age squared parameter shows a positive non-linear relationship between consumption levels and age.

The estimated parameters of the variables included in both hurdles have the expected signs and indicate that having a high education and being a white collar worker reduce the probability of smoking and the level of tobacco expenditure. The presence of children under fourteen years old significantly reduces the likelihood of smoking, indicating that households tend to refrain from smoking when small children are present. However, limited to the smokers sub-sample, the presence of children under fourteen years old increases the level of tobacco expenditure. The estimated coefficients for the dummy variable indicating the gender of the household's head (*MaleHH*) reveal

1  
2  
3  
4 the existence of differentiated behaviours, with female headed households presenting  
5 both a higher probability of smoking and higher tobacco expenditures. On the other  
6  
7 hand, the variable indicating the percentage of adult male members within the  
8  
9 household (*PercMale*) exerts a significant and positive effect on both participation and  
10  
11 consumption decisions. These evidences highlight the existence of important gender  
12  
13 differences in tobacco consumption patterns. Moreover, it is worth noticing that the  
14  
15 marital status variable exerts opposite effects on participation and consumption. In  
16  
17 particular, being a single reduces the likelihood to smoke but, conditional on smoking, it  
18  
19 increases tobacco expenditure.  
20  
21  
22  
23  
24

25  
26 Home ownership seems to negatively affect tobacco consumption levels; this result,  
27  
28 previously found by Jones (1989) and Yen (2005a), suggests that belonging to wealthier  
29  
30 social groups may induce moderate consumption.  
31  
32

33 Finally, the positive coefficient of the dummy variable indicating a high level of  
34  
35 alcohol expenditure (*HighAlc*) reveals that the probability of being a smoker is higher  
36  
37 for those households that are strong drinkers. Moreover, tobacco expenditure is also  
38  
39 found to be positively correlated with the level of alcohol expenditure, showing the  
40  
41 existence of a complementarity relationship in the consumption of addictive and habit-  
42  
43 generating goods.  
44  
45  
46  
47  
48

### 49 *5.3 Elasticities*

50  
51 In assessing the impact of explanatory variables, the presence of parameter estimates  
52  
53 with opposite signs in the two hurdles and the Box-Cox transformation complicate the  
54  
55 interpretation of the estimated effects. Thus, the impact of explanatory variables can be  
56  
57 better explored by computing elasticities.  
58  
59  
60



1  
2  
3  
4 The elasticities of probability, conditional level and unconditional level with respect  
5 to explanatory variables are calculated by using the formulas in equation (19). For  
6 statistical inference, standard errors are computed using the delta method (Su and Yen,  
7 1996; Spanos, 1999).  
8  
9  
10  
11  
12

13  
14  
15  
16 *(Table 5 about here)*  
17  
18  
19

20  
21 Estimated elasticities and discrete effects for the categorical variables, along with  
22 their standard errors, are presented in Table 5. Analysing the effects of the continuous  
23 variables, income has a positive effect on the conditional level of tobacco consumption;  
24 the elasticity are both significant, with a net effect on unconditional mean equal to  
25 0.2638, a value which is in line with the empirical findings of Yen (2005a). This finding  
26 shows that economic factors play an important role in determining smoking decision  
27 and indirectly suggest that observed zero consumption may well be the result not only  
28 of abstention but also of standard corner solution, confirming the validity of the double-  
29 hurdle specification for modelling tobacco expenditure.  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41

42 The elasticities with respect to age suggest that households with older heads are less  
43 likely to smoke and, conditional on smoking, consume less tobacco than younger  
44 households. This result is consistent with previous studies, even if the estimated effect  
45 on the unconditional level is lower than that found by Yen (1999).  
46  
47  
48  
49  
50

51 The level of alcohol consumption positively affects the unconditional level of  
52 tobacco consumption, revealing the existence of significant complementarities between  
53 smoking and drinking, with tobacco consumption rising as expenditures on alcoholic  
54 beverages grow.  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

The elasticities with respect to *PercMale* are all positive and significant, with an overall effect on the unconditional level equal to 0.0443. This result highlights how men and women exhibit different consumption patterns in relation to tobacco, with both smoking probability and conditional level increasing as the ratio of male to female members increases.

The analysis of the effects of binary variables gives further support to the evidence highlighted in the discussion of parameters estimates. In particular, it is worth noticing that education plays a negative and significant role only on the probability but not on the conditional level of smoking. The impact of education on the unconditional level of tobacco consumption is negative and significant at the 10 percent significance level (equal to -0.347) as the negative effect on probability obviously dominates. These evidences suggest that more educated individuals are more aware of the health risks associated with smoking and they are less likely to participate to tobacco consumption.

The occupational variable does not influence the probability of smoking, but white-collar workers are found to consume less tobacco conditional and unconditional on smoking.

The marital status variable exerts opposite effects on probability and conditional expenditure; more precisely, being a single reduces tobacco consumption probability by about 3 percent, but significantly increases the level of expenditure conditional on the smoking households sub-sample (the estimate effect is equal to 3.1230). However, given that the positive effect on conditional level dominates the negative effect on probability, the net effect on unconditional expenditure is clearly positive and equal to 1.3693.

Analysing the effect of household's head gender, it can be pointed out that the female-headed households are more likely to smoke and tend to consume more tobacco than the male-headed ones. The effects on probability, conditional and unconditional level are negative and significant, with an overall effect on the unconditional effect of equal to -0.7942. This result is not in contrast with the evidences connected to the effect of the percentage of male members and it is mainly connected to the structure of the female-headed households. In fact, the female-headed families in our sample are mainly single adult and single adult with adult children households and the effect exerted by this particular household structure may interact with the effect of household's head gender, making the separate identification of the two effects difficult.

Moreover, it should be noted that the categorical variable indicating the presence of children is characterized by a negative elasticity of probability (equal to -1.7%). However, conditional on smoking, the elasticity of consumption is positive and offsets the negative effect of probability.

Finally, heavy alcohol drinking households are characterized by positive and significant elasticities of probability, conditional and unconditional levels. In particular, this result suggests that the probability of smoking is higher (of about 5%) for those households displaying habit generation tendencies and less aware of the health risks connected with tobacco and alcohol consumption.

## 6. Concluding Remarks

In this paper a Box-Cox double-hurdle model has been applied to data on tobacco expenditure from the Italian Household Budget Survey. The test strategy to obtain the best specification that out-perform nested models was carried out by means of a

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

general-to-particular approach. The results obtained show that independence in the Box-Cox double-hurdle model is a non-restrictive assumption, while Tobit model, standard double-hurdle model with independence and Box-Cox Tobit model seem to be restrictive specifications, since they failed to account for non-normality in the bivariate distribution and for separate individual decisions on smoking participation and tobacco consumption.

The results based on the estimated parameters and elasticities confirm that decisions to smoke are related to income and demographic characteristics. In particular, income positively affects tobacco expenditure, indirectly showing the presence of habit-related behaviours in low income households which lead towards a lower income elasticity.

A substantial decline in the participation is found when the age of the household's head increases, together with an important lifecycle pattern in consumption. Estimation results highlight the presence of significant gender differences in both smoking participation and tobacco consumption patterns. High education and white collar occupation are found to reduce the likelihood of smoking and tobacco expenditure levels, suggesting that households with low educational levels and belonging to lower social classes have not yet benefit from policies concerning health consequences of smoking. Again, the data show that being a single determines a lower probability on smoking initiation, but conditional on smoking, the consumption level is higher.

Finally, complementarities between tobacco and alcoholic beverages consumption are present in both the probability and the level of smoking. This relationship is consistent with a wide body of empirical evidences and suggests that anti-smoking policies and public health strategies aimed at reducing alcohol abuse should be jointly addressed towards those households with higher levels of alcohol and tobacco expenditures.

1  
2  
3  
4  
5  
6  
7 **Acknowledgement.** We would like to thank Federico Perali and Luca Piccoli for their  
8 useful comments and suggestions. This paper is part of the research project “*Dynamic*  
9 *Analysis of Addiction: Intra-household Resources Allocation, Social Welfare and Public*  
10 *Health*”, University of Verona.  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20

## 21 **References**

- 22  
23 Amemiya, T. (1984) Tobit Models: A Survey, *Journal of Econometrics*, **84**, 3-61.  
24 Angulo, A.M., J.M. Gil and A. Gracia (2001) The demand for alcoholic beverages in  
25 Spain, *Agricultural Economics*, **26**, 71-83.  
26 Arabmazard, A. and P. Schmidt (1982) An Investigation of the Robustness of the Tobit  
27 Estimator to non-normality, *Econometrica*, **50**, 1055-1063  
28 Aristei, D., F. Perali and L. Pieroni (2005) Cohort Analysis of Alcohol Consumption: a  
29 Double-Hurdle Approach, Working Paper CHILD 09/2005.  
30 Atkinson, A.B., J. Gomulka, and N.H. Stern (1984) Household expenditure on tobacco  
31 1970-1980: evidence from the Family Expenditure Survey, ESRC Programme on  
32 Taxation, Incentives, and the Distribution of Income, London School of Economics,  
33 Discussion Paper No. 60.  
34 Bask, M. and M. Melkersson (2004) Rationally addicted to drinking and smoking?,  
35 *Applied Economics*, **36**, 373 – 381.  
36 Becker, G. S. and K. Murphy (1988) A Theory of Rational Addiction, *Journal of*  
37 *Political Economy*, **96**, 675-701.  
38 Becker, G.S., M. Grossman and K.M. Murphy (1994) An Empirical Analysis of  
39 Cigarette Addiction, *American Economic Review*, **84**, 396-418.  
40 Bera, A.K., C.M. Jarque, and L. Lee (1984) Testing the Normality Assumption in  
41 Limited Dependent Variable Models, *International Economic Review*, **25**, 563-78.  
42 Blaylock, J.R. and W.N. Blisard (1992) U.S. Cigarette consumption: The Case of Low-  
43 Income Women, *American Journal of Agricultural Economics*, **74**, 698-705.  
44 Blaylock, J.R. and W.N. Blisard (1993) Wine consumption by US men, *Applied*  
45 *Economics*, **24**, 645-651.  
46 Blundell, R. and C. Meghir (1987) Bivariate Alternatives to the Univariate Tobit Model,  
47 *Journal of Econometrics*, **34**, 179-200.  
48 Chaloupka, F. (1991) Rational Addictive Behavior and Cigarette Smoking, *Journal of*  
49 *Political Economy*, **99**, 722-42.  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3  
4  
5 Cragg, J. (1971) Some statistical models for limited dependent variables with  
6 application to the demand for durable goods, *Econometrica*, **39**, 829-844.
- 7  
8 Deaton, A. and M. Irish (1984) Statistical models for zero expenditures in household  
9 budgets, *Journal of Public Economics*, **23**, 59-80.
- 10  
11 Garcia, J. and J.M. Labeaga (1996) Alternative Approaches to Modelling Zero  
12 Expenditure: An Application to Spanish Demand for Tobacco, *Oxford Bulletin of*  
13 *Economics and Statistics*, **58**, 489-506.
- 14  
15 Heckman, J. (1979) Sample selection bias as a specification error, *Econometrica*, **47**,  
16 153-161.
- 17  
18 Jones, A.M. (1989) A double-hurdle model of cigarette consumption, *Journal of*  
19 *Applied Econometrics*, **4**, 23-39.
- 20  
21 Jones, A.M. (1992) A note on computation of the double-hurdle model with dependence  
22 with an application to tobacco expenditure, *Bulletin of Economic Research*, **44**, 67-74.
- 23  
24 Jones, A.M. and J.M. Labeaga (2003) Individual heterogeneity and censoring in panel  
25 data estimates of tobacco expenditure, *Journal of Applied Econometrics*, **18**, 157-  
26 177.
- 27  
28 Jones, A.M. and S.T. Yen (2000) A Box-Cox double-hurdle model, *The Manchester*  
29 *School*, **68**, 203-221.
- 30  
31 Kerr, W.C., T.K. Greenfield, J. Bond, Y. Ye, and Rehm, J. (2004) Age, period and  
32 cohort influences on beer, wine and spirits consumption trends in the US National  
33 Alcohol Surveys, *Addiction*, **99**, 1111-1120.
- 34  
35 Labeaga, J.M. (1999) A double-hurdle rational addiction model with heterogeneity:  
36 estimating the demand for tobacco, *Journal of Econometrics*, **93**, 49-72.
- 37  
38 Maddala, G.S. (1983) *Limited dependent and qualitative variables in econometrics*,  
39 Cambridge: Cambridge University Press.
- 40  
41 Maddala, G.S. and F.D. Nelson (1975) Specification Errors in Limited Dependent  
42 Variable Models, *NBER Working Paper 96*.
- 43  
44 McDonald, J. and R. Moffitt (1980) The Uses of Tobit Analysis, *The Review of*  
45 *Economics and Statistics*, **62**, 318-312.
- 46  
47 Moffatt, P.G. (2005) Hurdle models of loan default, *Journal of the Operational*  
48 *Research Society*, **56**, 1063-1071.
- 49  
50 Newman, C., M. Henchion, and A. Matthews (2003) A double-hurdle model of Irish  
51 household expenditure on prepared meals, *Applied Economics*, **35**, 1053-1061.
- 52  
53 Pagan, A. and P. Vella (1989) Diagnostic tests for models based on individual data: A  
54 survey, *Journal of Applied Econometrics*, **4**, S29-S59.
- 55  
56 Pudney, S. (1989) *Modelling Individual Choice: The Econometrics of Corners, Kinks*  
57 *and Holes*, New York: Basil Blackwell.
- 58  
59 Smith, M.D. (2003) On dependency in Double-Hurdle models. *Statistical Papers*, **44**,  
60 581-595.

- 1  
2  
3  
4  
5 Spanos, A. (1999) *Probability Theory and Statistical Inference: Econometric Modeling*  
6 *with Observational Data*. Cambridge, UK: Cambridge University Press.
- 7  
8 Su, S-J. and S.T. Yen (1996) Microeconomic Models of Infrequently Purchased  
9 Goods: An Application to Household Pork Consumption, *Empirical Economics*, **21**,  
10 513-533.
- 11  
12 Su, S-J. B. and S.T. Yen (2000) A censored system of cigarette and alcohol  
13 consumption, *Applied Economics*, **32**, 729-737.
- 14  
15 Tobin, J. (1958) Estimation of relationships for limited dependent variables,  
16 *Econometrica*, **26**, 24-36.
- 17  
18 Vuong, Q.H. (1989) Likelihood ratio test for model selection and non-nested  
19 hypotheses, *Econometrica*, **57**, 307-333.
- 20  
21 Wells, C. (2003) Retesting Fair's (1978) Model on Infidelity, *Journal of Applied*  
22 *Econometrics*, **18**, 237-239.
- 23  
24 Wooldridge, J.M. (2002) *Econometric Analysis of Cross Section and Panel Data*,  
25 Cambridge, MA: MIT Press.
- 26  
27 Yen, S.T. (1993) Working wives and food away from home: the Box-Cox double hurdle  
28 model, *American Journal of Agricultural Economics*, **75**, 884-895
- 29  
30 Yen, S.T. (1995) Alternative transformations in a class of limited dependent variable  
31 models: alcohol consumption by US women. *Applied Economics Letters*, **2**, 258-  
32 262.
- 33  
34 Yen, S.T. (1999) Gaussian versus count-data hurdle models: cigarette consumption by  
35 women in the US, *Applied Economics Letters*, **6**, 73-76.
- 36  
37 Yen, S.T. (2005a) A Multivariate Sample-Selection Model: Estimating Cigarette and  
38 Alcohol Demands with Zero Observations, *American Journal of Agricultural*  
39 *Economics*, **87**, 453-466.
- 40  
41 Yen, S.T. (2005b) Zero observations and gender differences in cigarette consumption,  
42 *Applied Economics*, **37**, 1839-1849
- 43  
44 Yen, S.T. and A.M. Jones (1996) Individual Cigarette Consumption and Addiction: a  
45 Flexible Limited Dependent Variable Approach, *Health Economics*, **5**, 105-117.
- 46  
47 Yen, S.T. and H.H. Jensen (1996) Determinants of household expenditures on alcohol,  
48 *The Journal of Consumer Affairs*, **30**, 48-67.
- 49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



## Tables

Table 1 – Variable definitions and descriptive statistics

Variable	Definition	Sample statistics			
		Full sample (n=27499)		Consuming (n=9184)	
		Mean	SD	Mean	SD
DEPENDENT VARIABLE					
TOBACCO	Per- adult household tobacco consumption	8.053	15.294	24.11264	17.696
EXPLANATORY VARIABLES (CONTINUOUS)					
AGE	Age of the household's head	56.057	15.818	51.730	13.634
AGESQR	Age of the household's head squared	3392.56	1822.87	2861.844	1482.522
PERCMALE	Percentage of adult male members in the household	0.456	0.272	0.510	0.237
INCOME	Proxied by per-equivalent adult household total expenditure and scaled by 100	11.166	8.580	11.946	9.019
INCOMESQR	Income squared	198.306	614.281	224.038	698.944
ALCOHOL	Per- adult household alcohol consumption	8.310	15.485	10.291	16.997
EXPLANATORY VARIABLES (BINARY)					
MALEHH	Equals 1 if the household's head is male	0.746		0.828	
HIGHEDU	Equals 1 if the household's head has at least a high school education, zero otherwise	0.286		0.294	
WHITECOLLAR	Equals 1 if the household's head is in a white collar occupation, zero otherwise	0.186		0.208	
OWNEROCC	Equals 1 if the household owns its home, zero otherwise	0.743		0.702	
SINGLE	Equals 1 for a single adult household without children, zero otherwise	0.290		0.207	
CHILD014	Equals 1 if any child aged 0-14 is present in the household, zero otherwise	0.243		0.290	
HIGHALC	Equals 1 if the household consumes a high level (over the 75 <sup>th</sup> percentile of the observed distribution) of alcoholic beverages, zero otherwise	0.555		0.658	



Table 2 – Diagnostic tests

Model	Homoscedasticity	Normality
<i>Tobit</i>	90.82 (2) [0.000]	134.603 (2) [0.000]
<i>Double-hurdle</i>	825.2 (3) [0.000]	138.122 (2) [0.000]

Note: the degrees of freedom of each  $\chi^2$  statistic are reported in round brackets while the p-value of each test is in squared brackets.

Table 3 – Specification tests

Model	Test type	Test value
<i>Box-Cox dependent double-hurdle vs. Box-Cox independent double-hurdle</i>	LR	0.78 (1) [0.3771]
<i>Box-Cox independent double-hurdle vs. Box-Cox Tobit</i>	Vuong	21.074*
<i>Box-Cox independent double-hurdle vs. Independent double-hurdle</i>	Vuong	38.428*
<i>Independent double-hurdle vs. Tobit</i>	Vuong	12.311*

Note: the degrees of freedom of the  $\chi^2$  statistic of the LR test are reported in round brackets while the corresponding p-value is in squared brackets. In the Vuong tests, the asterisk indicates that the null hypothesis of model equivalence is rejected at the 1% significance level.

Table 4 – Heteroscedastic double-hurdle estimates

Variable	Heteroscedastic Double-Hurdle Model			Box-Cox Heteroscedastic Double-Hurdle Model		
	Participation	Consumption	Het.	Participation	Consumption	Het.
AGE	-0.0032 (5.60)	-0.9120 (7.08)	–	-0.0101 (16.42)	-0.0560 (9.02)	–
AGESQR	–	0.0067 (5.66)	–	–	0.0004 (7.84)	–
MALEHH	-0.0453 (4.48)	-2.2069 (2.26)	–	-0.0704 (6.52)	-0.0884 (2.13)	–
HIGHEDU	-0.0354 (4.43)	0.0886 (0.14)	–	-0.0327 (3.91)	-0.0297 (0.98)	–
WHITECOLLAR	-0.0035 (0.39)	-2.5902 (3.48)	–	-0.0182 (1.78)	-0.1666 (4.86)	–
PERCMALE	0.1764 (8.49)	3.2854 (2.59)	–	0.2461 (11.93)	0.1646 (2.66)	–
SINGLE	-0.1071 (8.57)	23.5328 (33.75)	–	-0.0804 (8.28)	1.1662 (30.94)	–
CHILD014	-0.0451 (4.36)	1.6787 (2.36)	–	-0.0435 (3.87)	0.1237 (3.89)	–
OWNEROcc	–	-4.6780 (8.07)	–	–	-0.2355 (8.46)	–
INCOME	–	12.8827 (8.05)	0.2489 (15.00)	–	0.6113 (14.34)	0.0476 (2.77)
INCOMESQR	–	-3.1280 (6.22)	-0.0078 (6.55)	–	-0.0819 (8.75)	0.0047 (1.97)
ALCOHOL	–	0.0994 (2.84)	0.005 (5.49)	–	0.0077 (9.40)	–
HIGHALC	0.1014 (8.51)	–	–	0.1309 (11.67)	–	–
CONSTANT	0.0261 (1.21)	27.1526 (7.57)	–	0.6647 (16.70)	4.5645 (26.40)	–
Box-Cox parameter ( $\lambda$ )					0.1864 (17.50)	
Log Likelihood		-50881.02			-31097.36	

Note: the absolute value of each t-statistic is reported in round brackets.

Table 5 – Elasticities with respect to continuous variables  
and effects of binary variables

<i>Variables</i>	<i>Probability</i>	<i>Conditional level</i>	<i>Unconditional level</i>
<i>Continuous variables</i>			
AGE	-0.2551*** (0.0155)	-0.0748*** (0.0040)	-0.3299*** (0.0134)
PERCMALE	0.0339*** (0.0042)	0.0104*** (0.0022)	0.0443*** (0.0011)
INCOME	–	0.2638*** (0.0184)	0.2638*** (0.0184)
ALCOHOL	–	0.0025*** (0.0003)	0.0025*** (0.0003)
<i>Discrete variables</i>			
MALEHH	-0.0272*** (0.0104)	-0.8969*** (0.0561)	-0.7942*** (0.0426)
HIGHEDU	-0.0127*** (0.0058)	-0.2952 (0.2460)	-0.3470* (0.2015)
WHITECOLLAR	-0.0071 (0.0088)	-1.4820*** (0.4940)	-0.5296** (0.2493)
SINGLE	-0.0313*** (0.0094)	3.1230*** (0.0919)	1.3693*** (0.0437)
CHILD014	-0.0169** (0.0085)	1.2554*** (0.0523)	0.7603*** (0.0335)
OWNEROcc	0.0462*** (0.0067)	-2.1037*** (0.0915)	-0.5305*** (0.0286)
HIGHALC	0.0506*** (0.0062)	2.0469*** (0.1279)	1.0921*** (0.0341)

Notes: Asymptotic standard errors of estimated elasticities and discrete effects are reported in round brackets. Asterisks indicate levels of significance: \*\*\* = 0.01, \*\* = 0.05 and \* = 0.10.

1  
2  
3  
4  
5 **Figures**  
6  
7  
8  
9  
10

11 *Figure 1 – Nested Models*  
12

