

## A randomised controlled trial of cognitive behavioural treatment for obsessive compulsive disorder in children and adolescents

Williams, Tim I.; Salkovskis, Paul M.; Forrester, Liz; Turner, Sam; White, Hilary; Allsopp, Mark A.

Postprint / Postprint

Zeitschriftenartikel / journal article

Zur Verfügung gestellt in Kooperation mit / provided in cooperation with:

[www.peerproject.eu](http://www.peerproject.eu)

### Empfohlene Zitierung / Suggested Citation:

Williams, T. I., Salkovskis, P. M., Forrester, L., Turner, S., White, H., & Allsopp, M. A. (2009). A randomised controlled trial of cognitive behavioural treatment for obsessive compulsive disorder in children and adolescents. *European Child & Adolescent Psychiatry*, 19(5), 449-456. <https://doi.org/10.1007/s00787-009-0077-9>

### Nutzungsbedingungen:

Dieser Text wird unter dem "PEER Licence Agreement zur Verfügung" gestellt. Nähere Auskünfte zum PEER-Projekt finden Sie hier: <http://www.peerproject.eu> Gewährt wird ein nicht exklusives, nicht übertragbares, persönliches und beschränktes Recht auf Nutzung dieses Dokuments. Dieses Dokument ist ausschließlich für den persönlichen, nicht-kommerziellen Gebrauch bestimmt. Auf sämtlichen Kopien dieses Dokuments müssen alle Urheberrechtshinweise und sonstigen Hinweise auf gesetzlichen Schutz beibehalten werden. Sie dürfen dieses Dokument nicht in irgendeiner Weise abändern, noch dürfen Sie dieses Dokument für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, aufführen, vertreiben oder anderweitig nutzen.

Mit der Verwendung dieses Dokuments erkennen Sie die Nutzungsbedingungen an.

**gesis**  
Leibniz-Institut  
für Sozialwissenschaften

### Terms of use:

This document is made available under the "PEER Licence Agreement". For more information regarding the PEER-project see: <http://www.peerproject.eu> This document is solely intended for your personal, non-commercial use. All of the copies of this documents must retain all copyright information and other information regarding legal protection. You are not allowed to alter this document in any way, to copy it for public or commercial purposes, to exhibit the document in public, to perform, distribute or otherwise use the document in public.

By using this particular document, you accept the above-stated conditions of use.

Mitglied der  
  
Leibniz-Gemeinschaft

# A randomised controlled trial of cognitive behavioural treatment for obsessive compulsive disorder in children and adolescents

Tim I. Williams · Paul M. Salkovskis · Liz Forrester ·  
Sam Turner · Hilary White · Mark A. Allsopp

Received: 4 November 2008 / Accepted: 4 November 2009 / Published online: 17 November 2009  
© Springer-Verlag 2009

**Abstract** Cognitive behaviour therapy (CBT) for young people with obsessive compulsive disorder (OCD) has become the treatment of first choice. However, the literature is largely based on studies emphasising exposure and response prevention. In this study, we report on a randomised controlled trial of CBT for young people carried out in typical outpatient clinic conditions which focused on cognitions. A randomised controlled trial compares 10 sessions of manualised cognitive behavioural treatment with a 12-week waiting list for adolescents and children with OCD. Assessors were blind to treatment allocation. 21 consecutive patients with OCD aged between 9 and 18 years were recruited. The group who received treatment improved more than a comparison group who waited for 3 months. The second group was treated subsequently using the same protocol and made similar gains. In conclusion, CBT can be delivered effectively to young people with OCD in typical outpatient settings.

**Keywords** Obsessive compulsive disorder · Cognitive behaviour therapy · Randomised controlled trial · Treatment · Young people

## Introduction

Obsessive compulsive disorder (OCD) is particularly a severe and disabling psychological disorder. Recent epidemiological studies suggest that it affects around 1.9–3.2% of the adult population and may have a 1-year prevalence rate of up to 4% in late adolescence [13]. Follow-up studies have shown that it has a chronic relapsing course such that 50% of adult patients report their first symptoms in childhood or adolescence and 50% of patients with OCD in adolescence will continue to suffer disabling effects from OCD in adulthood [2, 8, 35]. An effective treatment for OCD in adolescence could offer significant savings for the health service and improved quality of life for many patients.

For adults, pharmacological treatments (selective serotonin re-uptake inhibitors, SSRIs) have been shown to reduce the level of symptomatology, but in up to 90% of patients these gains are lost within 7 weeks of stopping treatment [34]. The benefits of SSRIs for young people with OCD have also been demonstrated (see [17] for a meta-analysis), but there are concerns about the safety of medication in young people on the grounds of both physiology and risk taking behaviour. Although controversial, these concerns impact on the acceptability of pharmacotherapy. In adults, it is known that SSRIs do not enhance short-term adherence to psychological treatment, and in the long term their use may impair the efficacy of psychological treatments [15].

Recent expert consensus guidelines suggest that cognitive behavioural treatments are the first choice treatments for children and adolescents with OCD [21, 26] although a Cochrane review concluded only that “behavioural or cognitive behaviour therapy (CBT) appears to be a promising treatment for OCD in children

---

T. I. Williams (✉)  
Berkshire Healthcare NHS Trust, School of Psychology,  
University of Reading, Earley Gate, Reading RG6 6AL, UK  
e-mail: sxswiams@reading.ac.uk

P. M. Salkovskis · L. Forrester  
Department of Psychology, Institute of Psychiatry,  
Kings College, University of London, London, UK

S. Turner · H. White · M. A. Allsopp  
Berkshire Adolescent Service, Berkshire Healthcare NHS Trust,  
Wokingham Hospital, Wokingham, UK

and adolescents” [27]. Until recently, there have been few trials which examine the effectiveness of cognitive behavioural therapy for young people, based on the assumption that adult-based research would generalise. In general, controlled trials have demonstrated that CBT techniques used for adults have generalised well [4, 6, 9, 11, 22]. The largest of these trials [22] demonstrated that both medication and cognitive behavioural therapy were effective treatments for OCD. However, there was a significant site  $\times$  treatment interaction such that in one site CBT was significantly more effective than medication whereas in the other site medication was found to be more effective. Other controlled trials of CBT for young people have found similar beneficial effects of CBT compared with medication [4, 11].

The components of CBT for children have often been poorly specified. Two components have been used by most studies—exposure and response prevention (E/RP) and anxiety management [4, 6, 9, 11, 22]. Bolton and Perrin [9] have demonstrated that E/RP alone (with minimal anxiety management) is sufficient to achieve significant benefits. De Haan et al. [11] investigated the use of targeted cognitive techniques including manipulating responsibility cognitions demonstrating comparable effects for CBT and medication. Two studies have delivered CBT in group format [4, 6], without compromising the effect of CBT. Furthermore, Barrett et al. [6] involved parents in treatment with beneficial effects.

An alternative to managing anxiety around exposure and response prevention is to target cognitions specific to OCD. A number of candidate cognitive processes have been suggested and a systematic review of cognitions related to OCD symptoms in young people [28] suggested that three models have some support: increased responsibility, meta-cognitions, and thought–action fusion. These models are difficult to distinguish empirically, because the processes overlap. An appraisal of responsibility for intrusive thoughts is a form of meta-cognition (thinking about thoughts) and the appraisal that one’s thoughts might result in actions or events occurring is of no significance unless one appraises oneself as responsible for the thought. Evidence for responsibility cognitions being specific to OCD in adults has been provided by Salkovskis and his co-workers [31] who demonstrated that they differentiate patients with OCD from those with other emotional disorders.

Studies with young people have not shown the clarity of the link between responsibility and OCD expected from the adult results [28]. Other aspects of cognition such as thought–action fusion [7, 19, 23] and perfectionism [19] are correlated with symptoms of OCD in young people, although these relationships are substantially mediated by responsibility cognitions [19, 24].

Reynolds and Reeves [28] point out, however, that the relationships between these cognitive processes are unclear and may be mediated or moderated by other factors such as low mood. The robustness of the responsibility cognitions model in adults and the evidence that responsibility cognitions seem to be problematic in children with OCD suggest that there is utility in examining the extent to which CBT which concentrated on responsibility cognitions could be used to treat OCD.

Salkovskis and his colleagues [29] have developed a package of cognitive behavioural treatment methods which concentrates on responsibility cognitions. The components that seem to be particularly important are:

1. normalising the nature of unpleasant intrusive cognitions;
2. identifying the nature of the link between the thoughts and the feelings of discomfort/anxiety and subsequent neutralising rituals;
3. examining the logical nature of the links, and putting them to the test by means of behavioural experiments;
4. comparing real danger with the worry about causing harm, i.e. discriminating between thinking about and acting upon;
5. helping the patient to identify that the effect of attempting to control the thoughts leads to recurrence of the thoughts or images;
6. helping the patient to consider alternative non-threatening accounts of their obsessive problems.


The therapy also includes psychoeducation about anxiety and an explanation of how trying to carry out exposure and response prevention might seem difficult. Although some exposure work was undertaken, it was always explained in terms of finding out what happens to cognitions and emotions. The therapists did not insist on waiting until the uncomfortable feelings had subsided as is required by exposure and response prevention. A pilot study [36] of this form of CBT with young people suggested that the changes in symptoms were paralleled by the changes in the responsibility beliefs which initiate and maintain compulsive behaviours. The techniques also affect meta-cognition in that they alter the beliefs about the nature and meaning of cognitions associated with OCD. This form of CBT differs from that used in the POTS trial [22] in its concentration on cognitions. The treatment does not aim to teach the children to resist the impulse to carry out compulsions, but rather to identify and change the misconceptions underlying the motivation to carry out compulsions. A much larger study would be needed to test this form of CBT against other forms of CBT or indeed a purely behavioural treatment such as that in Bolton and Perrin [9].

**Methods**

Young people were recruited to the trial with a primary diagnosis of OCD based on a semi-structured interview for mental health problems (ADIS-C) [33]. In total, 21 young people aged 9–18 years (mean age 13 years 7 months; 13 boys, 8 girls) were recruited from 22 cases referred by Child and Adolescent Mental Health Teams or Family Doctors for the trial (see Fig. 1 for CONSORT diagram). Children were included if OCD was the major problem, and it had been present for at least 6 months. Children were excluded if they were unable to speak and understand English fluently, if they had co-occurring psychosis or autism spectrum disorder. At initial assessments, 11 presented with no other clinical diagnoses, while 4 received diagnoses of generalised anxiety disorder, 4 specific

phobia, 4 separation anxiety, 2 ADHD, 2 social phobia, and 1 dysthymia. (The numbers do not add up to 10 because only 2 had one additional diagnosis, the others having up to three additional diagnoses.) Seven of the participants were taking medication throughout the trial and had been taking the same dose for 12 weeks prior to the trial (two on paroxetine 5 mg/day, three on fluoxetine 20 mg/day, one on fluvoxamine 50 mg/day, one on clomipramine 50 mg/day). There was no significant multivariate difference between the treated group and the waiting list group at initial assessment [ $F(6, 13) = 0.58, n.s.$ ]. The therapists for the trial (TW, HW, ST) were clinical psychologists employed by the National Health Service (NHS) in England to work in community child and adolescent mental health clinics. All treatment took place in NHS clinics and was recorded on audiotape if the young person consented.

**Fig. 1** CONSORT diagram



## Procedure

Once consent was received for taking part in the trial, the participants were allocated to either immediate treatment (ten 1-h sessions) or a 12-week waiting list. Allocation was carried out on a predetermined random number schedule with no replacements by the trial administrator. Participants in the waiting list condition were informed that they would have to wait 12 weeks for therapy, but they were given the phone number of the lead clinician (TW) to contact in the case of a significant deterioration in OCD.

Assessments were conducted at the beginning and end of treatment (and/or waiting list period), and 12 weeks after the end of treatment. All assessments were completed in the child's home unless specifically requested to be elsewhere. Assessors were blind to the allocation of the participants, and the participants were instructed not to reveal whether they had received treatment. The assessors completed two semi-structured interviews: Children's Yale-Brown Obsessive Compulsive Scale (C-YBOCS) [32] to assess OCD symptoms, and the ADIS-C [33], a semi-structured interview to assess the presence of other comorbid disorders. Participant completed measures included Child Depression Inventory [18], Obsessions and Compulsions Inventory (OCI) [14] modified for children, Multidimensional Anxiety Scale for Children [20], the Children's Responsibility Attributions Scale (CRAS), the Children's Responsibility Interpretations questionnaire (CRIQ). Both the CRIQ and the CRAS are instruments modified from the adult measures of responsibility cognitions published in Salkovskis et al. [31]. The CRAS is scored such that increasing score indicates a decreasing level of responsibility attributions, whereas the other self-report scales are scored such that increasing score indicates increasing difficulties. An unpublished study by the first author found that the internal reliability of the CRAS was good in a large normative sample of 13–14-year olds (Cronbach's  $\alpha = 0.85$ ). The internal reliability in a sample of young people with mental health problems for the other scales was also high (OCI: Cronbach's  $\alpha = 0.93$ ; RIQ frequency: Cronbach's  $\alpha = 0.86$ ; RIQ belief: Cronbach's  $\alpha = 0.85$ ). Copies of these measures are available on <http://psychology.iop.kcl.ac.uk/ocdkids/questionnaires/questionnaires.aspx>.

## Treatments

Cognitive behaviour therapy was based on the principles outlined by Salkovskis [29]. Participants worked with their therapists to understand the cognitive distortions which maintain their OCD. The aim of treatment is to alter responsibility cognitions, primarily by doing experiments both in session and at home. For instance, one common

belief encountered in OCD is that the sufferer is uniquely responsible for harm occurring if certain rituals are not carried out satisfactorily. The therapist and the participant would agree during the session to carry out an experiment designed to see what happens if responsibility is shared. Another common task during therapy is to attempt to elicit the worrying intrusive thoughts and examine whether they have real meaning or are just thoughts. Treatment fidelity was ensured through clinic notes, regular meetings of the therapist team and audiotapes of treatment sessions.

Participants were allocated by the trial administrator to the two groups using a table of random numbers. Only the trial administrator was aware which participants were in which group.

## Results

### Plan of analysis

The primary measure of OCD symptoms is the CYBOCS. We first report an analysis of covariance of the CYBOCS score at 3-month post-baseline assessment (with baseline CYBOCS as the covariate), at which stage only one of the two groups will have received an active treatment. This analysis tests the hypothesis that there will be a group effect at 3 months. We will then repeat the analysis of covariance of the CYBOCS scores at 6-month post-baseline when both groups will have received treatment. This analysis tests the hypothesis that there will be a period by group effect on the analysis of the CYBOCS scores at 3 and 6 months. Finally, we will analyse the secondary measures (i.e. self-report data) in the same way using initial values of the measure as the covariate.

Two clients dropped out of the study during the first phase (one from each condition). In the following analyses, the last observation was carried forward from the last available observation. This is, therefore, an intention to treat analysis.

### Analysis of primary measure


Figure 2 shows the mean CYBOCS scores at baseline, 3 and 6 months. The analysis of covariance (baseline CYBOCS as covariate) of the CYBOCS at 3 months showed a significant group effect [ $F(1) = 7.07$ ,  $p = 0.016$ ]. The figure shows that this is due to a much improved CYBOCS score for the group treated with CBT first. The group that was allocated to the waiting list showed little or no improvement over the same period. Cohen's effect size ( $d$ ) for the difference between the two groups divided by the mean standard deviation was calculated as 1.07, which is considered to be a large treatment effect.



At 6 months, the analysis of covariance of CYBOCS (using baseline CYBOCS as covariate) showed no group effect as expected since both groups had received treatment, although there was a trend towards a significant effect of the covariate [ $F(1, 19) = 3.70, p = 0.07$ ] suggesting that the severity of OCD at the beginning of the trial influences the outcome. As can be seen from Fig. 2, both groups had improved significantly and the group that was treated first showed continuing slight improvement.

#### Analysis of secondary measures

Analysis of the secondary data (the self-report questionnaires OCI, CRAS, CDI, CRIQ frequency, CRIQ belief; see Table 1) used the same procedures, i.e. analysis of covariance using the baseline value of the measure as the covariate. There were no statistically significant group effects at 3 months [OCI:  $F(1, 19) = 0.29, p = 0.59$ ; CDI:  $F(1, 19) = 0.49, p = 0.49$ ; CRIQ frequency:  $F(1, 19) = 2.07, p = 0.17$ ; CRIQ belief:  $F(1, 19) = 0.90, p = 0.36$ ; CRAS:  $F(1, 19) = 0.41, p = 0.53$ ; MASC:  $F(1, 19) = 0.008, p = 0.929$ ] but there were significant effects of the covariate for each measure [OCI:  $F(1, 19) = 11.19, p = 0.004$ ; CDI:  $F(1, 19) = 12.20, p = 0.003$ ; CRIQ frequency:  $F(1, 19) = 13.06, p = 0.002$ ; CRIQ belief:  $F(1, 19) = 17.26, p = 0.001$ ; CRAS:  $F(1, 19) = 27.12, p < 0.001$ ], except the MASC [ $F(1, 19) = 0.13, p = 0.72$ ]. Inspection of Table 2 shows that the groups were improving on all measures at the 3-month point.



**Fig. 2** Mean CYBOCS scores for both groups at baseline, 3 and 6 months. Error bars represent standard errors of the mean at each time point

A second analysis of covariance was carried out to determine if the treatment was effective for both groups using data from baseline and 6 months later after both groups had received CBT. The results showed that there was only a significant time effect [ $F(1, 19) = 53.14, p < 0.001$ ] and no group [ $F(1, 19) = 0.06, p = 0.82$ ] or interaction effect [ $F(1, 19) = 0.73, p = 0.40$ ]. As shown in Fig. 1, the changes in CYBOCS are largest when the groups are receiving treatment. Overall, the CYBOCS scores of the participants reduced from 22.12 (SE = 1.08) to 9.64 (SE = 1.79). Analysis of the secondary data also showed significant changes from baseline to 6 months [OCI:  $F(1, 19) = 18.35, p < 0.001$ ; CDI:  $F(1, 19) = 23.16, p < 0.001$ ; RIQB:  $F(1, 19) = 32.8, p < 0.001$ ; RIQF:  $F(1, 19) = 25.15, p < 0.001$ ; RAS:  $F(1, 19) = 10.75, p = 0.004$ ; MASC:  $F(1, 19) = 22.81, p < 0.001$ ], but no group effects [OCI:  $F(1, 19) = 0.45, p = 0.51$ ; CDI:  $F(1, 19) = 0.53, p = 0.48$ ; RIQB:  $F(1, 19) = 0.30, p = 0.59$ ; RIQF:  $F(1, 19) = 0.003, p = 0.38$ ; RAS:  $F(1, 19) = 0.89, p = 0.36$ ; MASC:  $F(1, 19) = 0.30, p = 0.60$ ] or interaction effects [OCI:  $F(1, 19) = 1.41, p = 0.25$ ; CDI:  $F(1, 19) = 0.42, p = 0.53$ ; RIQB:  $F(1, 19) = 1.03, p = 0.32$ ; RIQF:  $F(1, 19) = 0.83, p = 0.38$ ; RAS:  $F(1, 19) = 2.08, p = 0.17$ ; MASC:  $F(1, 19) = 0.27, p = 0.61$ ]. Table 1 shows the means and standard deviations for all the measures.

We followed the recommendations of Morris and DeShon (2002 cited in [1]) to calculate the effect size as the difference between the immediately pre-treatment CYBOCS and the post-treatment CYBOCS divided by the pre-treatment standard deviation (effect size = 2.62) which also allows for the calculation of the effect size of the waiting list. This effect size was compared with those from other published trials which included a control group (see Table 2). The results from this study fall within the range of values reported heretofore.

#### Discussion

Cognitive behaviour therapy largely based on the use of experiments to tackle the cognitive biases in OCD produced a significantly greater reduction in OCD symptoms of the participants than a waiting list condition. Being placed on the waiting list first did not affect the power of the subsequent treatment. The self-report measures showed a statistically significant reduction in self-reported symptoms over the two time periods but did not demonstrate an effect of treatment. The study also shows only small changes in symptoms for the young people placed on the waiting list (cf. [1]), therefore confirming the chronic nature of OCD in young people. This adds to the body of evidence in favour of CBT for OCD in young people. In

**Table 1** Mean (SE) of self-report measures at baseline, 3 months (after first phase, one group treated) and 6 months (after second phase, both groups treated)

Measure	Baseline		3 months		6 months	
	CBT	WL	CBT	WL	CBT	WL
CYBOCS	23.09 (1.22)	21.05 (1.84)	12.09 (2.25)	19.60 (2.03)	9.23 (2.45)	10.10 (2.74)
OCI	59.30 (8.28)	73.55 (8.26)	45.00 (8.30)	60.30 (9.62)	37.10 (8.73)	34.30 (5.51)
CDI	17.85 (2.76)	14.67 (1.82)	12.9 (2.62)	12.78 (2.92)	10.50 (2.41)	9.06 (2.56)
MASC	59.8 (6.87)	66.3 (6.74)	49.7 (5.90)	56.6 (6.69)	41.1 (4.40)	43.0 (7.28)
CRAS	49.00 (6.00)	51.00 (8.10)	58.00 (7.54)	63.63 (6.15)	58.00 (8.03)	74.13 (8.31)
CRIQ belief	634.38 (103.66)	781.11 (152.50)	563.13 (99.26)	537.78 (150.76)	320.63 (86.94)	332.22 (102.34)
CRIQ frequency	29.25 (4.31)	31.44 (5.14)	25.38 (3.90)	19.33 (5.45)	16.75 (4.36)	13.89 (4.35)

Gallop [16] reports means of 64.17 for CRAS (higher scores indicate less responsibility attributions), 451.7 for CRIQ belief and 17.4 for frequency (where higher scores indicate more responsibility interpretations)

**Table 2** Effect sizes on CYBOCS calculated according to Morris and DeShon [25] for controlled studies of CBT in young people

Study	Individual CBT effect size	Waiting list or placebo control effect size
This study	-2.61792	-0.24914
March et al. [22] (placebo)	-2.6087	-0.81818
De Haan et al. [11]	-2.10169	N/A
Barrett et al. [6]	-3.55349	0.198543

Negative numbers indicate improved CYBOCS scores. N/A indicates that the study did not include a waiting list control

view of the potential problems associated with medication, the results of this study support the view that young people with OCD should, therefore, be offered CBT as the first-line treatment by child and adolescent mental health services.

This is the first randomised controlled study of a CBT approach based on responsibility cognitions (following [29, 30]) with young people. Although the numbers treated in this trial were small, the demonstration that the treatment was as effective for the waiting list group demonstrates that waiting for treatment had no effect on eventual outcome and increases the power of the study. The 3-month follow-up of the group that was treated first suggests that, at least at first, the participants continued to improve after cessation of treatment.

The failure of the self-report measures to demonstrate a difference between the groups at the 3-month point is difficult to explain. Although a failure to find changes in depression symptoms is not unusual in trials of CBT for OCD [25], we had predicted a change in OCD symptoms paralleling responsibility cognitions as was found in a pilot study [36]. One possibility is that as found by Anholt et al. [3] the changes in behaviour rather than the changes in cognitions were the most significant feature of treatment.

However, it is also possible that the hope that someone else was going to help with the problem improved the subjective feelings tapped by self-report measures including the appraisals of responsibility or affected the responsibility cognitions directly (e.g. someone else is taking charge of the OCD for me), but did not alter the assessor-rated severity. If that is so, it might indicate that changing cognitions alone are not sufficient to alter behaviour. Indeed, the individual formulations of OCD used in this treatment method use multiple maintenance cycles which are described for each affected person. Yet another possibility is that the self-report measures are not valid in this population. Further research on the validity and reliability of self-report measures would be helpful.

Other trials of CBT with young people have tended to concentrate on managing the anxiety or discomfort experienced when undertaking exposure and response prevention, e.g. [4, 6, 22]. The effect size observed in this trial is somewhat less than those seen in previous trials of CBT for young people with OCD (mean effect size 1.98, Table 3 in [1]), although the differences in the methods of effect size calculation make the comparison somewhat problematic.

Differences in the treatment offered may also be important in determining the effectiveness of the intervention. There are two lines of evidence: intra-trial site differences and inter-trial differences. The POTS trial compared CBT using E/RP and management of the distress associated with OCD with a placebo medication condition but found considerable site differences for both the CBT and a third medication-only condition [22]. (Unfortunately because the site data are only reported as Hedge's *g* effect sizes, we were unable to compare effect sizes using the Morris and DeShon method [25] and calculated Hedge's *g* effect sizes in the following text.) The Hedge's *g* effect size for the POTS trial form of CBT varied from 0.51 in Chapel Hill to 1.6 in Pennsylvania (a statistically significant

difference). The Hedge's  $g$  effect size in this trial (1.07) falls within that range. Bolton and Perrin [9] demonstrated the benefits of E/RP alone compared with a wait list (Hedge's  $g$  effect size 1.23).

Our study does not provide information on the relative merits of different forms of CBT. The difficulty with attempting such a comparison is that the effect sizes of different models of CBT appear to be comparable (see previous paragraph). Of more interest in clinical practice are issues such as the use of combinations of medication and CBT and the involvement of the family. Studies with adults have begun to show that offering medication first diminishes the subsequent effectiveness of CBT [15].

Most of the studies on CBT with young people have included the parents in treatment sessions, although Bolton and Perrin [9] only provided feedback at the end of each session. Recent work by Barrett et al. [6] showed that very substantial reductions in OCD could be obtained by the use of family management components in the treatment package. The families of young people with OCD differ in some important respects from the families of young people with anxiety disorders [5, 12]. Barrett's study [5] demonstrated that the parents were less confident in their child's ability, less rewarding of independence, and were less likely to use positive problem solving than the parents of children with anxiety disorders. Derisley et al. [12] found that parents tended to use avoidant coping techniques as well as having more symptoms of mental health problems. The NICE guidelines too [26] suggest that the contribution of the family to the treatment of young people with OCD needs further investigation. However, the involvement of families in the treatment of other emotional disorders in young people has not always resulted in more improvement than a less explicit component of family change [10]. Nevertheless, it would be worth investigating whether a family component focused on problem-solving strategies and developing confidence in the child's abilities (based on Barrett's studies [5, 6]), and would be helpful in increasing the effect of the CBT used in this study as has been partially demonstrated by Barrett et al. [6].

**Acknowledgments** This research was supported by a grant (SPGS 808) from the National Health Service Responsive Funding Scheme to the first author.

## References

- Abramowitz JS, Whiteside SR, Deacon BJ (2005) The effectiveness of treatment for pediatric obsessive-compulsive disorder: a meta-analysis. *Behav Ther* 36(1):55–63
- Allsopp M, Verduyn C (1989) A follow-up of adolescents with obsessive-compulsive disorder. *Br J Psychiatry* 154:829–834
- Anholt GE, Kempe P, de Haan E, van Oppen P, Cath DC, Smit JH et al (2008) Cognitive versus behavior therapy: processes of change in the treatment of obsessive-compulsive disorder. *Psychother Psychosom* 77(1):38–42
- Asbahr FR, Castillo AR, Ito LM, Latorre MD, Moreira MN, Lotufo-Neto F (2005) Group cognitive-behavioral therapy versus sertraline for the treatment of children and adolescents with obsessive-compulsive disorder. *J Am Acad Child Adolesc Psychiatry* 44(11):1128–1136
- Barrett P, Shortt A, Healy L (2002) Do parent and child behaviours differentiate families whose children have obsessive-compulsive disorder from other clinic and non-clinic families? *J Child Psychol Psychiatry* 43:597–607
- Barrett P, Healy-Farrell L, March JS (2004) Cognitive-behavioral family treatment of childhood obsessive-compulsive disorder: a controlled trial. *J Am Acad Child Adolesc Psychiatry* 43:46–62
- Barrett P, Healy-Farrell L (2003) Perceived responsibility in juvenile obsessive-compulsive disorder: an experimental manipulation. *J Clin Child Adolesc Psychol* 32:430–441
- Bolton D, Luckie M, Steinberg D (1995) Long-term course of obsessive-compulsive disorder treated in adolescence. *J Am Acad Child Adolesc Psychiatry* 34:1441–1450
- Bolton D, Perrin S (2008) Evaluation of exposure with response-prevention for obsessive compulsive disorder in childhood and adolescence. *J Behav Ther Exp Psychiatry* 39:11–22
- Bodden DHM, Bögels SM, Nauta MH, De Haan E, Ringrose J, Appelboom C et al (2008) Child versus family cognitive-behavioral therapy in clinically anxious youth: an efficacy and partial effectiveness study. *J Am Acad Child Adolesc Psychiatry* 47(12):1384–1394
- De Haan E, Hoogduin KAL, Buitelaar JK, Keijsers GPJ (1998) Behavior therapy versus clomipramine for the treatment of obsessive-compulsive disorder in children and adolescents. *J Am Acad Child Adolesc Psychiatry* 37:1022–1029
- Derisley J, Libby S, Clark S, Reynolds S (2005) Mental health, coping and family-functioning in parents of young people with obsessive-compulsive disorder and with anxiety disorders. *Br J Clin Psychol* 44:439–444
- Douglass HM, Moffitt TE, Dar R, McGee R et al (1995) Obsessive-compulsive disorder in a birth cohort of 18-year-olds: prevalence and predictors. *J Am Acad Child Adolesc Psychiatry* 34:1424–1431
- Foa EB, Kozak MJ, Salkovskis PM, Coles ME, Amir N (1998) The validation of a new obsessive-compulsive disorder scale: The Obsessive-Compulsive Inventory. *Psychol Assess* 10:206–214
- Foa EB, Liebowitz MR, Kozak MJ, Davies S, Campeas R, Franklin ME, Huppert JD, Kjemristed K, Rowan V, Schmidt AB, Simpson HB, Tu X (2005) Randomized, placebo-controlled trial of exposure and ritual prevention, clomipramine, and their combination in the treatment of obsessive-compulsive disorder. *Am J Psychiatry* 162:151–161
- Gallop C (2009) Cognitive behavioural assessment of OCD. In: Waite PL, Williams TI (eds) *Obsessive compulsive disorder: cognitive behaviour therapy with children and young people*. Routledge, London
- Geller DA, Biederman J, Stewart SE, Mullin B, Martin A, Spencer T, Faraone SV (2003) Which SSRI? A meta-analysis of pharmacotherapy trials in pediatric obsessive-compulsive disorder. *Am J Psychiatry* 160:1919–1928
- Kovacs M (1992) *The children's depression inventory manual*. Multi-Health Systems, Toronto
- Libby S, Reynolds S, Derisley J, Clark S (2004) Cognitive appraisals in young people with obsessive-compulsive disorder. *J Child Psychol Psychiatry* 45:1076–1084
- March JS (1997) *Technical manual for the multidimensional anxiety scale for children (MASC)*. Multi-Health Systems, New York



21. March JS, Frances A, Carpenter D, Kahn MD (1997) Treatment of obsessive-compulsive disorder. The Expert Consensus Panel for obsessive-compulsive disorder. *J Clin Psychiatry* 58(Suppl 4):2–72
22. March JS, Foa E, Gammon P, Chrisman A, Curry J, Fitzgerald D, Sullivan K, Franklin M, Huppert J, Rynn M, Zhao N, Zoellner L, Leonard H, Garcia A, Freeman J, Tu X (2004) Cognitive-behavior therapy, sertraline, and their combination for children and adolescents with obsessive-compulsive disorder—the Pediatric OCD Treatment Study (POTS) randomized controlled trial. *J Am Med Assoc* 292:1969–1976
23. Mather A, Cartwright-Hatton S (2004) Cognitive predictors of obsessive-compulsive symptoms in adolescence: a preliminary investigation. *J Clin Child Adolesc Psychol* 33:743–749
24. Matthews L, Reynolds S, Derisley J (2006) Examining cognitive models of obsessive compulsive disorder in adolescents. *Behav Cogn Psychother* 35(02):149–163
25. Morris SB, DeShon RP (2002) Combining effect size estimates in meta-analysis with repeated measures and independent-groups designs. *Psychol Methods* 7:105–125
26. National Collaborating Centre for Mental Health (2006) Obsessive compulsive disorder: core interventions in the treatment of obsessive compulsive disorder and body dysmorphic disorder. National Clinical Practice Guideline Number 31. The British Psychological Society, The Royal College of Psychiatrists, London
27. O’Kearney RT, Anstey KJ, von Sanden C (2006) Behavioural and cognitive behavioural therapy for obsessive compulsive disorder in children and adolescents. *Cochrane Database Syst Rev*, no. 4, art. no. CD004856. doi:[10.1002/14651858.CD004856.pub2](https://doi.org/10.1002/14651858.CD004856.pub2)
28. Reynolds S, Reeves J (2008) Do cognitive models of obsessive compulsive disorder apply to children and adolescents? *Behav Cogn Psychother* 36(04):463–471
29. Salkovskis PM (1998) Psychological approaches to the understanding of obsessional problems. In: Swinson RP, Antony MM, Rachman SJ, Richter MA (eds) *Obsessive-compulsive disorder: theory, research and treatment*. Guilford, New York
30. Salkovskis PM (1999) Understanding and treating obsessive-compulsive disorder. *Behav Res Ther* 37:s29–s52
31. Salkovskis P, Wroe A, Gledhill A, Morrison N, Forrester E, Richards C, Reynolds M, Thorpe S (2000) Responsibility attitudes and interpretations are characteristic of obsessive compulsive disorder. *Behav Res Ther* 38:347–372
32. Scahill L, Riddle MA, McSwiggin Hardin M, Ort SI, King RA, Goodman WK, Cicchetti D, Leckman JF (1997) Children’s Yale-Brown Obsessive Compulsive Scale: reliability and validity. *J Am Acad Child Adolesc Psychiatry* 36:844–852
33. Silverman WK, Albano AM (1996) Anxiety disorders interview schedule for DSM-IV: child and parent versions. The Psychological Corporation, San Antonio
34. Simpson HB, Franklin ME, Cheng JF, Foa EB, Liebowitz MR (2005) Standard criteria for relapse are needed in obsessive-compulsive disorder. *Depression Anxiety* 21:1–8
35. Thomsen PH, Mikkelsen HU (1995) Course of obsessive compulsive disorder in children and adolescents. A prospective follow-up study of 23 Danish cases. *J Am Acad Child Adolesc Psychiatry* 34:1432–1440
36. Williams T, Salkovskis P, Forrester EA, Allsopp MA (2002) Changes in symptoms of OCD and appraisal of responsibility during cognitive behavioural treatment: a pilot study. *Behav Cogn Psychother* 30:69–78